



# The efficacy of platelet gel derived from umbilical cord blood on diabetic foot ulcers: A double-blind randomized clinical trial

Seyedeh Esmat Hosseini<sup>b,g,1</sup>, Behnam Molavi<sup>a,1</sup>, Alireza Goodarzi<sup>c</sup>, Ahad Alizadeh<sup>d</sup>,  
Alireza Yousefzadeh<sup>e</sup>, Niloofar Sodeifi<sup>f</sup>, Leila Arab<sup>g</sup>, Nasser Aghdami<sup>g,\*</sup>

<sup>a</sup> Sina Trauma and Surgery Research Center, Tehran University of Medical Sciences, Tehran, Iran

<sup>b</sup> Student Research Committee, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>c</sup> Department of Medical Laboratory Sciences, School of Paramedicine, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>d</sup> Metabolic Diseases Research Center, Research Institute for Prevention of Non-Communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran

<sup>e</sup> Department of Epidemiology and Reproductive Health, Reproductive Epidemiology Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

<sup>f</sup> Department of Andrology at Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

<sup>g</sup> Department of Regenerative Medicine, Stem Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

## ARTICLE INFO

### Keywords:

Diabetic foot ulcer  
Platelet gel  
Platelet-derived growth factor  
Umbilical cord blood

## ABSTRACT

**Introduction:** Type 2 diabetes is one of the most prevalent diseases throughout the world. The foot ulcers are severe complications of this disease. Foot ulcers induce the high rate of morbidity, impair quality of life, bring about extreme costs to health service providers, and give rise to waste of time. Recently, platelet-rich plasma (PRP) and platelet gel (PG) have been used for the treatment of chronic wounds. In the present randomized, double-blind, placebo-controlled study, platelet gel derived from umbilical cord blood (UCB) was used to heal the diabetic foot.

**Method:** The patients were randomly divided into three groups, under the categories of PG, platelet-poor plasma (PPP) and lubricant gel (placebo) (ratio 1:1:1). The processes of gels application were launched for the subject of each group twice per week with 3–4 days' interval. This mechanism protracted for eight weeks. After completion of 8 weeks, the patients were followed up after two weeks and then continued for nine months with one-month interval.

**Result:** 30 patients underwent statistical analysis. Except for diastolic blood pressure which was significant between groups, there were no statistically significant differences in patients' baseline demographics. No significant differences were detected between groups at baseline of wounds ( $P = 0.09$ ). The results revealed that there is no statistically significant interaction among three groups during follow-up time.

**Conclusion:** The present study provides evidence that there are no significant differences in the size of wound among PG, PPP, and placebo groups.

## 1. Introduction

Type 2 diabetes are considered worldwide incidence and prevalence [1]. According to the World Health Organization (WHO), 366 million people will have been affected diabetic by 2030 [2]. Approximately 12–25 % of these afflicted patients are at risk of ulcer particularly in the extremities in their life span [2–4]. The lower limb ulcers are diabetes serious complications [3]. Foot ulcers cause significant morbidity, impair quality of life, incur high costs to health service providers and time

-consuming process [4–6]. These wounds can lead to the patients' hospitalization and may result in amputation of limbs [7]. Since, the wound healing is a slow process, it often takes between 2 and 5 months and requires special treatments [8]. Holzer et al. in a retrospective analysis of costs for lower extremity ulcers in people with diabetes come to this conclusion that new treatment strategies should be developed owing to the wound treatment high medical cost [4]. Various interventional approaches have been used for the treatment of diabetes-induced ulcer. One of the treatment methods of the chronic wounds is

\* Corresponding author at: Tehran Province, Tehran, Masoud St., Iran.

E-mail addresses: [esmat.hosseini\\_110@yahoo.com](mailto:esmat.hosseini_110@yahoo.com) (S.E. Hosseini), [molavibe@sina.tums.ac.ir](mailto:molavibe@sina.tums.ac.ir) (B. Molavi), [agoodarzi58@yahoo.com](mailto:agoodarzi58@yahoo.com) (A. Goodarzi), [st.alizadeh@gmail.com](mailto:st.alizadeh@gmail.com) (A. Alizadeh), [alireza.yousefzadekh@gmail.com](mailto:alireza.yousefzadekh@gmail.com) (A. Yousefzadeh), [sniloofar@yahoo.com](mailto:sniloofar@yahoo.com) (N. Sodeifi), [leara91@gmail.com](mailto:leara91@gmail.com) (L. Arab), [nasseragh@yahoo.com](mailto:nasseragh@yahoo.com) (N. Aghdami).

<sup>1</sup> These authors contributed equally to this work.

<https://doi.org/10.1016/j.wndm.2020.100178>

Received 2 January 2019; Received in revised form 9 December 2019; Accepted 30 December 2019

Available online 08 January 2020

2213-9095/ © 2020 Published by Elsevier GmbH.